

AD-A209 351

OFFICE OF NAVAL RESEARCH

Contract N00014-86-K-0043

TECHNICAL REPORT No. 101

Electron Energy Levels in a Quantum Well Within an In-Plane Magnetic Field

by

H. R. Lee, H. G. Oh, Thomas F. George and C. I. Um

Prepared for Publication

in

Journal of Applied Physics

Departments of Chemistry and Physics  
State University of New York at Buffalo  
Buffalo, New York 14260

June 1989

Reproduction in whole or in part is permitted for any purpose of the  
United States Government.

This document has been approved for public release and sale;  
its distribution is unlimited.

DTIC  
ELECTE  
JUN 20 1989  
S E D

89 6 19 124

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE

## REPORT DOCUMENTATION PAGE

Form Approved  
OMB No. 0704-0188

1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION / AVAILABILITY OF REPORT Approved for public release; distribution unlimited		
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S)  UBUFFALO/DC/89/TR-101			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Depts. Chemistry & Physics State University of New York		6b. OFFICE SYMBOL (If applicable)		7a. NAME OF MONITORING ORGANIZATION	
6c. ADDRESS (City, State, and ZIP Code) Fronczak Hall, Amherst Campus Buffalo, New York 14260		7b. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217			
8a. NAME OF FUNDING / SPONSORING ORGANIZATION Office of Naval Research		8b. OFFICE SYMBOL (If applicable)		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER  Contract N00014-86-K-0043	
8c. ADDRESS (City, State, and ZIP Code) Chemistry Program 800 N. Quincy Street Arlington, Virginia 22217		10. SOURCE OF FUNDING NUMBERS			
		PROGRAM ELEMENT NO.		PROJECT NO.	
		TASK NO.		WORK UNIT ACCESSION NO.	
11. TITLE (Include Security Classification)  Electron Energy Levels in a Quantum Well Within an In-Plane Magnetic Field					
12. PERSONAL AUTHOR(S) H. R. Lee, H. G. Oh, Thomas F. George and C. I. Um					
13a. TYPE OF REPORT		13b. TIME COVERED FROM _____ TO _____		14. DATE OF REPORT (Year, Month, Day) June 1989	
				15. PAGE COUNT 17	
16. SUPPLEMENTARY NOTATION  Prepared for publication in Journal of Applied Physics					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	ELECTRON ENERGY LEVELS		
			CYCLOTRON ORBITS		
			QUANTUM WELL		
			QUANTUM CALCULATION,		
			IN-PLANE MAGNETIC FIELD		
			BULK LANDAU LEVELS, (1e)		
19. ABSTRACT (Continue on reverse if necessary and identify by block number)					
<p>The exact eigenenergy spectrum of an electron is calculated in a quantum well within an in-plane magnetic field. The numerical solutions for the excited energy states as well as the ground state energy are found for various quantum-well widths and barrier heights. The cyclotron orbits are considerably affected by the quantum well. The energy levels higher than the potential height of the quantum well and the energy levels lower than the potential height of the quantum well show quite different behaviors. These are explained with the properties of the combined potential and the wave function inherent to the system.</p>					
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Dr. David L. Nelson			22b. TELEPHONE (Include Area Code) (202) 696-4410		22c. OFFICE SYMBOL

Electron Energy Levels in a Quantum Well Within an In-Plane Magnetic Field

H. R. Lee, H. G. Oh and Thomas F. George  
Department of Physics and Astronomy  
239 Fronczak Hall  
State University of New York at Buffalo  
Buffalo, New York 14260

C. I. Um  
Department of Physics  
College of Science  
Korea University  
Seoul 136, KOREA



Abstract

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

The exact eigenenergy spectrum of an electron is calculated in a quantum well within an in-plane magnetic field. The numerical solutions for the excited energy states as well as the ground state energy are found for various quantum-well widths and barrier heights. The cyclotron orbits are considerably affected by the quantum well. The energy levels higher than the potential height of the quantum well and the energy levels lower than the potential height of the quantum well show quite different behaviors. These are explained with the properties of the combined potential and the wave function inherent to the system.

PACS Nos.: 73.60.-n, 73.20.Dx, 71.50.+t

## I. Introduction

The properties of electrons confined in semiconductor heterostructures have been studied for different situations. The carriers in two-dimensional systems subjected to a magnetic field perpendicular to the layers are completely quantized into Landau levels, which have been extensively studied with respect to the quantized Hall effect[1] and Shubnikov-de-Haas oscillations [2]. In this case, since the magnetic field is in the same direction as the confining electric field, the Hamiltonian can be separated into an electric part leading to subbands and a magnetic part leading to Landau levels. For any other orientation, this separation is not possible any more. Thus, in an external magnetic field parallel to the interface, the situation becomes more complicated. By studying the effects of magnetic field on the optical transitions, we may obtain the details of the subband structure[3,4]. Although an in-plane magnetic field usually has little effect on two-dimensional properties, it can strongly affect the spectrum of intersubband optical transitions[5]. If the magnetic length  $a_H = (\hbar/\mu\omega)^{1/2}$ , where  $\mu$  and  $\omega$  are the effective mass and the cyclotron frequency of an electron, respectively, is made comparable to the quantum well width by increasing the magnitude of the magnetic field, the problem is even more complicated because the confining electric field and the magnetic field contribute almost the same weight to the energy levels of electrons. There are several papers in theory[6-9] as well as in experiment[10-12] which study the effect of in-plane magnetic fields on two-dimensional systems. Recently Klama[8] has reported the quantization rules and analytic expressions of the electron energy spectrum in a thin film within a

longitudinal magnetic field. In his paper, the electronic motion is always inside the quantum well because the potential height is taken to be infinite. The eigenenergies of two-dimensional electrons subjected to a tilted magnetic field are solved analytically in a parabolic potential well[10], which is useful for the analysis of transport measurements in their experiments. Oliveira et al[9] have found the electron binding energy in a quantum well of n-type GaAs-Al<sub>x</sub>Ga<sub>1-x</sub>As heterostructures by a self-consistent solution of the Poisson and Schrödinger equations. Most of their calculations were done for a couple of the lowest subbands in various systems.

Since the harmonic oscillator is one of the exactly solvable problems[13], the exact eigenvalues for a quantum well under an in-plane magnetic field can be obtained numerically. In this paper we present the numerical solutions for the energy spectrum of electrons in a quantum well with a finite potential height  $V_0$  and well width  $W$  comparable to the magnetic length. The eigenenergy spectrum of an infinite potential well is also calculated for comparison. In the calculations, the normalized potential height  $(V_0/\hbar\omega)$  and well width  $(\sqrt{2}W/a_H)$  are used to see the results for an arbitrary potential height  $V_0$  and well width  $W$ .

## II. Theory

Let us consider a quantum well with well width  $W$  and the potential height  $V_0$  in an external applied longitudinal DC magnetic field. The uniform external magnetic field  $\vec{B} = (0, 0, B_0)$  oriented in the plane of the layer can be described by the vector potential  $\vec{A} = (0, B_0x, 0)$ . In the direction parallel to  $\vec{B}$  the electrons have a free motion, whereas in the

plane perpendicular to  $\vec{B}$  they are subjected to the combined potential of the magnetic field and the quantum well. Then, the Hamiltonian for an electron in a potential well with a magnetic field in the  $xz$ -plane can be written as

$$H = \frac{1}{2\mu} (p_x^2 + p_z^2) + \frac{1}{2\mu} \left( p_y + eB_0 x \right)^2 + V(x) \quad (1)$$

where  $\mu$  is the effective mass and  $V(x)$  is the potential energy of the electron in the well. Correspondingly, the Schrödinger equation is given by

$$H\psi(\vec{r}) = E\psi(\vec{r}) \quad (2)$$

Due to the translational symmetry in the  $y$ - and  $z$ -directions, the wave function can be written as

$$\psi(\vec{r}) = \chi(x) e^{i(q_y y + q_z z)} \quad (3)$$

Substituting this wave function into the Schrödinger equation (2), we obtain the effective one-dimensional Schrödinger equation

$$\frac{d^2 \chi(x)}{dx^2} + \frac{2\mu}{\hbar^2} \left( \epsilon - \frac{\mu \omega^2}{2} (x - x_0)^2 + V(x) \right) \chi(x) = 0 \quad (4)$$

where

$$\epsilon = E - \frac{p_z^2}{2\mu}, \quad (5)$$

and we have used  $\omega = eB_0/\mu$  for the cyclotron frequency,  $x_0 = a_H^2 q_y$  for the position of the orbit center and  $a_H = (\hbar/\mu\omega)^{1/2}$  for the magnetic length.

Now the two boundaries caused by the quantum well can be determined depending on the location of the orbit center  $x_0$ . For the general case, we assume the left and right boundaries of the well are located at  $x_L = a$  and  $x_R = b$  ( $a < b$ ), respectively, i.e.,

$$V(x) = \begin{cases} 0 & \text{for } a < x < b \\ V_0 & \text{for } x < a, \quad b < x \end{cases} \quad (6)$$

Introducing the dimensionless length scale  $\xi = \sqrt{2}x/a_H$  for the  $x$ -coordinate, the Schrödinger equation for the combined potential is divided into three regions:

$$\frac{d^2 x_I(\xi)}{d\xi^2} + \left[ \frac{\epsilon}{\hbar\omega} - \frac{1}{4} (\xi - \xi_0)^2 \right] x_I(\xi) = 0 \quad \text{for } \frac{\sqrt{2}a}{a_H} < \xi < \frac{\sqrt{2}b}{a_H} \quad (7a)$$

$$\frac{d^2 x_{II}(\xi)}{d\xi^2} + \left[ \frac{\epsilon - V_0}{\hbar\omega} - \frac{1}{4} (\xi - \xi_0)^2 \right] x_{II}(\xi) = 0 \quad \text{for } \xi < \frac{\sqrt{2}a}{a_H} \quad (7b)$$

$$\frac{d^2 x_{III}(\xi)}{d\xi^2} + \left[ \frac{\epsilon - V_0}{\hbar\omega} - \frac{1}{4} (\xi - \xi_0)^2 \right] x_{III}(\xi) = 0 \quad \text{for } \frac{\sqrt{2}b}{a_H} < \xi \quad (7c)$$

Comparing the above Schrödinger equations (7a), (7b), and (7c) with the Weber equation [14],

$$\frac{d^2\psi(z)}{dz^2} + \left(m + \frac{1}{2} - \frac{1}{4}z^2\right)\psi(z) = 0, \quad (8)$$

we can identify the eigenvalues of the system:

$$\frac{\epsilon}{\hbar\omega} = m + \frac{1}{2} \quad \text{for} \quad \frac{\sqrt{2a}}{a_H} < \xi < \frac{\sqrt{2b}}{a_H} \quad (9)$$

and

$$\frac{\epsilon - V_0}{\hbar\omega} = m' + \frac{1}{2} \quad \text{for} \quad \xi < \frac{\sqrt{2a}}{a_H}, \text{ or } \frac{\sqrt{2b}}{a_H} < \xi. \quad (10)$$

The quantum numbers  $m$  and  $m'$  are related each other by,

$$(m - m')\hbar\omega = V_0. \quad (11)$$

The solution of equation (8) is the well-known Weber function[14]:

$$D_m(z) = 2^{m/2} e^{-z^2/4} \left[ \frac{\sqrt{\pi}}{\Gamma(1/2 - m/2)} F\left(-\frac{1}{2}m \middle| \frac{1}{2} \middle| \frac{1}{2}z^2\right) - \frac{\sqrt{2\pi} z}{\Gamma(-m/2)} F\left(\frac{1}{2} - \frac{1}{2}m \middle| \frac{3}{2} \middle| \frac{1}{2}z^2\right) \right] \quad (12)$$

where  $F(a|b|x)$  is the confluent hypergeometric function[15] and  $\Gamma(z)$  is the gamma function.

From the asymptotic properties of the Weber function[14], the general solutions of the Schrödinger equations (7a), (7b) and (7c) may take the following forms:



$$x_I(\xi) = A D_m(\xi) + B D_m(-\xi) \quad (13a)$$

$$x_{II}(\xi) = C D_{m'}(-\xi) \quad (13b)$$

$$x_{III}(\xi) = F D_{m'}(\xi) \quad (13c)$$

where A, B, C and F are the normalization constants in each region. Then the continuity of the logarithmic derivatives of the wave functions at both boundaries  $\xi_L = \sqrt{2}a/a_H$  and  $\xi_R = \sqrt{2}b/a_H$  of the quantum well provides the following equation for the energy eigenvalues:

$$\begin{aligned} & D_m(\xi_R) D'_m(\xi_R) - D'_{m'}(\xi_R) D_m(\xi_R) \\ & - \frac{D'_m(\xi_L) D_{m'}(-\xi_L) + D'_{m'}(-\xi_L) D_m(\xi_L)}{D'_m(-\xi_L) D_{m'}(-\xi_L) + D'_{m'}(-\xi_L) D_m(-\xi_L)} \left[ D'_{m'}(\xi_R) D_m(-\xi_R) + D_{m'}(\xi_R) D'_m(-\xi_R) \right] = 0 \end{aligned} \quad (14)$$

where

$$D'_m(\xi) = \frac{d}{d\xi} D_m(\xi) \quad (15)$$

### III. Results and Discussion

Making use of the recursion relations of the Weber function[14], the numerical solutions for m can be obtained from Eq. (14) if the values of  $V_0$ ,

$\xi_L$  and  $\xi_R$  are given. The numerical solutions of  $m$  are illustrated in Figs. (1) and (2) for different values of normalized potential height  $V_0/\hbar\omega$  and well width  $W/a_H$ . We use the potential height  $V_0 = 5\hbar\omega$  and the well width  $W = a_H/\sqrt{2}$  in Fig. 1 and  $V_0 = 10\hbar\omega$  and  $W = \sqrt{2}a_H$  were used in Fig. 2. In these figures, the various energy levels are plotted against the position  $\xi_0$  of the orbit center. In each graph the dotted line shows the boundaries of the quantum well on a normalized scale, and the quantum number  $m$  (y-axis) is related to the eigenenergy by  $\epsilon = (m + 1/2)\hbar\omega$ . These results clearly show two different types of energy states: (1) the states confined in the quantum well (the lowest level in Fig. 1 and the lowest two levels in Fig. 2) and (2) the extended states (the higher levels in Figs. 1 and 2).

For the confined states (1), most of the wavefunction is pushed into the quantum well because the eigenenergy is lower than the potential height. As the orbit center is shifted from the center of the potential well, the electron penetrates and spends increasingly more time inside the potential barrier and less time in the well, resulting thus in an increase of energy. If the orbit center is moved deep into the barrier and far away from the quantum well, the low-level electrons see essentially a magnetic harmonic well elevated by  $V_0$ . This means that the energy versus  $\xi_0$  curve will flatten out at large values of  $\xi_0$ . Hence the usual bulk Landau levels take over except for a shift of  $V_0$  in energy. Before the bulk Landau levels take over, we can see the crossing of the confined state and the extended states in the intermediate shift of orbit center  $\xi_0$  in both Figs. 1 and 2. These two states are degenerate at the moment they cross each other. If the orbit center is shifted into the barrier, the quantum well would be elevated as

well as changed in shape by the presence of the parabolic magnetic potential. This means that the overall enhancement of the quantum well due to the magnetic potential should be parabolic as the orbit center  $\xi_0$  is moving away from the quantum well. Thus the eigenenergies of the confined states in the quantum well increase parabolically as a function of the orbit center  $\xi_0$ . Clearly the crossings of the confined states and the extended states in Figs.1 and 2 exhibit parabolic behavior. The changes in shape of the quantum well neglecting the elevation have little effect on the crossings because the eigenenergies of the confined states are always limited by the barrier height  $V_0$ , which is small compared to the elevation due to the magnetic potential.

In the second case (2), the energy spacing between neighboring modes in the energy spectrum is about  $\hbar\omega$ , but each mode makes oscillations as the orbit center moves away from the center of the quantum well. In this case, the electron energies are higher than the potential height  $V_0$  and the wavefunctions are located partly within the well and partly inside the barrier, but still confined by the magnetic potential well. The oscillatory behavior of the eigenmodes seems to be related to the oscillating property of the harmonic wavefunction. The harmonic wavefunction  $[D_m(z)]$  has many nodes and peaks depending on the quantum number  $m$ , and the wavefunction may be classified as symmetric (even parity) or antisymmetric (odd parity) for the centered orbit ( $\xi_0=0$ ). The oscillations of eigenenergy in each mode depend on the number of nodes on the wave function. Thus the number of oscillations increases one by one as we go up to the higher levels. We also observe that the oscillations of the eigenenergy are within the range of the

wavepacket size ( $\Delta x = \sqrt{m+1/2} a_H$ ). The alternating downward and upward concavity in energy spectrum at the middle of the quantum well seems due to the symmetric or antisymmetric properties of the wavefunction. If the wavefunction is symmetric, the middle peak of the wavefunction is located at the center of the potential well for the centered orbit ( $\xi_0=0$ ). Therefore the eigenenergy will increase if the orbit center is shifted slightly toward barrier, resulting in the downward concavity. On the other hand, we expect upward concavity for the antisymmetric wavefunction because its node is now located at the lowest spot of the combined potential well. If these arguments are correct, the wavefunction of the lowest mode should be symmetric and have no node, the second mode should be antisymmetric and have one node, and so on from Figs. 1 and 2. In conclusion, we can say that the energy levels with oscillating structures are direct images of the magnetic wave function itself in the restricted geometry.

For the practical example of  $\text{GaAs-Al}_x\text{GaAs}_{1-x}$  with  $x = 0.3$ , we get  $V_0 = 0.25$  eV. To match this potential height in fig.1 where  $V_0 = 5\hbar\omega$ , we need  $B = 29$  T, which corresponds to  $a_H = 48$  Å. If  $V_0 = 10\hbar\omega$  like in Fig. 2, a magnetic field of  $B = 14$  T is required. To observe these results at  $B = 10$  T which has a magnetic length  $a_H = 81$  Å, the potential height of Figs. 1 and 2 correspond to  $V_0 = 87$  and 174 meV, respectively. Even if these results were calculated in a relatively strong magnetic field due to the computer limit, the general features could be observed in a practical samples given above.

Finally, the eigenenergies for an infinite potential well ( $V_0 \rightarrow \infty$ ) can be calculated similarly with the proper boundary conditions ( $\chi_{II} = \chi_{III} = 0$ ,

$\chi_I(\xi - \xi_R) - \chi_I(\xi - \xi_L) = 0$ ) and the results are presented in Fig. 3 with a well width  $W = \sqrt{2}a_H$ . In this case, since all the electronic motions are within the quantum well, we cannot see such extended states as for a finite potential height. As explained earlier, the parabolic behavior is almost the same except for higher energies than before due to the infinite well.

#### Acknowledgments

This research was supported by the Office of Naval Research and the Office of Air Force Office of Scientific Research (AFSC), United States Air Force, under Contract F49620-86-C-0009. The United States Government is authorized to reproduce and distribute reprints for governmental purposes, notwithstanding any copyright notation hereon.

# References

- [1] E. von Klitzing, G. Dorda, and M. Pepper, Phys. Rev. Lett. 45, 494 (1980), D. C. Tsui, H. L. Störmer, and A. C. Gossard, Phys. Rev. Lett. 48, 1559 (1982), and R. B. Laughlin, Phys. Rev. B, 23, 5632 (1981).
- [2] A. B. Fowler, F. F. Fang, W. E. Howard and P. J. Stiles, Phys. Rev. Lett. 16, 901(1966), and K. F. Komatsubara, K. Narita, Y. Katayama, N. Kotera, and M. Kobayashi, J. Phys. Chem. Solids 35, 723 (1974).
- [3] W. Beinvogl and J. F. Koch, Phys. Rev. Lett. 40, 1736 (1978).
- [4] F. Koch, in Physics in High Magnetic Fields, edited by S. Chikazumi and M. Miura (Springer, Berlin, 1981).
- [5] T. Ando, A. B. Fowler, and F. Stern, Rev. Mod. Phys. 54, 437 (1982).
- [6] T. Ando, J. Phys. Soc. Japan 39, 411 (1975).
- [7] S. K. Bhattacharya, Phys. Rev. B 25, 3756 (1982).
- [8] S. Klama, J. Phys. C: Solid State Phys. 20, 551 (1987).
- [9] Glaucia M. G. Oliveira, Vivili M. S. Gomes, A. S. Chaves, J. R. Leite and J. M. Worlock, Phys. Rev. B 35, 2896 (1987).
- [10] J. C. Maan, in Two-dimensional Systems. Heterostructures and Superlattices, Vol.53 in Solid State Sciences, edited by G. Bauer, F. Kuchar and H. Heinrich, (Springer-Verlag, Berlin, 1984).
- [11] J. H. Caseman and U. Merkt, Solid State Commun. 47, 917 (1983).
- [12] Z. Schlesinger, J. C. M. Hwang and S. J. Allen Jr., Phys. Rev. Lett. 50, 2098 (1983)
- [13] E. Schrödinger, Naturwirs 14, 664 (1926).
- [14] P. M. Morse and H. Feshbach, Methods of Theoretical Physics (McGraw-Hill, New York, 1953), p. 1403.
- [15] M. Abramowitz and I. A. Stegun, Handbook of Mathematical Functions, (Dover, New York, 1964), p. 503.

Figure Captions

1. Numerical solution for electron eigenenergy as a function of the orbit center  $\xi_0$  with well width  $W = a_H/\sqrt{2}$  and potential height  $V_0 = 5\hbar\omega$ .
2. Numerical solution for electron eigenenergy as a function of the orbit center  $\xi_0$  with well width  $W = \sqrt{2}/a_H$  and potential height  $V_0 = 10\hbar\omega$ .
3. Numerical solution for electron eigenenergy as a function of the orbit center  $\xi_0$  with well width  $W = \sqrt{2}/a_H$  and potential height  $V_0 = \infty$ .

Fig. 1






Fig. 2





Fig. 3



TECHNICAL REPORT DISTRIBUTION LIST, GEN

	<u>No. Copies</u>		<u>No. Copies</u>
Office of Naval Research Attn: Code 1113 800 N. Quincy Street Arlington, Virginia 22217-5000	2	Dr. David Young Code 334 NORDA NSTL, Mississippi 39529	1
Dr. Bernard Douda Naval Weapons Support Center Code 50C Crane, Indiana 47522-5050	1	Naval Weapons Center Attn: Dr. Ron Atkins Chemistry Division China Lake, California 93555	1
Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko, Code L52 Port Hueneme, California 93401	1	Scientific Advisor Commandant of the Marine Corps Code RD-1 Washington, D.C. 20380	1
Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	12 high quality	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 12211 Research Triangle Park, NC 27709	1
DTNSRDC Attn: Dr. H. Singerman Applied Chemistry Division Annapolis, Maryland 21401	1	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112	1
Dr. William Tolles Superintendent Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375-5000	1	Naval Ocean Systems Center Attn: Dr. S. Yamamoto Marine Sciences Division San Diego, California 91232	1
		Dr. David L. Nelson Chemistry Division Office of Naval Research 800 North Quincy Street Arlington, Virginia 22217	1

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. J. E. Jensen  
Hughes Research Laboratory  
3011 Malibu Canyon Road  
Malibu, California 90265

Dr. C. B. Harris  
Department of Chemistry  
University of California  
Berkeley, California 94720

Dr. J. H. Weaver  
Department of Chemical Engineering  
and Materials Science  
University of Minnesota  
Minneapolis, Minnesota 55455

Dr. F. Kutzler  
Department of Chemistry  
Box 5055  
Tennessee Technological University  
Cookeville, Tennessee 38501

Dr. A. Reisman  
Microelectronics Center of North Carolina  
Research Triangle Park, North Carolina  
27709

Dr. D. DiLella  
Chemistry Department  
George Washington University  
Washington D.C. 20052

Dr. M. Grunze  
Laboratory for Surface Science and  
Technology  
University of Maine  
Orono, Maine 04469

Dr. R. Reeves  
Chemistry Department  
Renssaele Polytechnic Institute  
Troy, New York 12181

Dr. J. Butler  
Naval Research Laboratory  
Code 6115  
Washington D.C. 20375-5000

Dr. Steven M. George  
Stanford University  
Department of Chemistry  
Stanford, CA 94305

Dr. L. Interante  
Chemistry Department  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. Mark Johnson  
Yale University  
Department of Chemistry  
New Haven, CT 06511-8118

Dr. Irvin Heard  
Chemistry and Physics Department  
Lincoln University  
Lincoln University, Pennsylvania 19352

Dr. W. Knauer  
Hughes Research Laboratory  
3011 Malibu Canyon Road  
Malibu, California 90265

Dr. K.J. Klaubunde  
Department of Chemistry  
Kansas State University  
Manhattan, Kansas 66506

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. G. A. Somorjai  
Department of Chemistry  
University of California  
Berkeley, California 94720

Dr. J. Murday  
Naval Research Laboratory  
Code 6170  
Washington, D.C. 20375-5000

Dr. J. B. Hudson  
Materials Division  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. Theodore E. Madey  
Surface Chemistry Section  
Department of Commerce  
National Bureau of Standards  
Washington, D.C. 20234

Dr. J. E. Demuth  
IBM Corporation  
Thomas J. Watson Research Center  
P.O. Box 218  
Yorktown Heights, New York 10598

Dr. M. G. Lagally  
Department of Metallurgical  
and Mining Engineering  
University of Wisconsin  
Madison, Wisconsin 53706

Dr. R. P. Van Duyne  
Chemistry Department  
Northwestern University  
Evanston, Illinois 60637

Dr. J. M. White  
Department of Chemistry  
University of Texas  
Austin, Texas 78712

Dr. D. E. Harrison  
Department of Physics  
Naval Postgraduate School  
Monterey, California 93940

Dr. R. L. Park  
Director, Center of Materials  
Research  
University of Maryland  
College Park, Maryland 20742

Dr. W. T. Peria  
Electrical Engineering Department  
University of Minnesota  
Minneapolis, Minnesota 55455

Dr. Keith H. Johnson  
Department of Metallurgy and  
Materials Science  
Massachusetts Institute of Technology  
Cambridge, Massachusetts 02139

Dr. S. Sibener  
Department of Chemistry  
James Franck Institute  
5640 Ellis Avenue  
Chicago, Illinois 60637

Dr. Arnold Green  
Quantum Surface Dynamics Branch  
Code 3817  
Naval Weapons Center  
China Lake, California 93555

Dr. A. Wold  
Department of Chemistry  
Brown University  
Providence, Rhode Island 02912

Dr. S. L. Bernasek  
Department of Chemistry  
Princeton University  
Princeton, New Jersey 08544

Dr. W. Kohn  
Department of Physics  
University of California, San Diego  
La Jolla, California 92037

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. F. Carter  
Code 6170  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. Richard Colton  
Code 6170  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. Dan Pierce  
National Bureau of Standards  
Optical Physics Division  
Washington, D.C. 20234

Dr. R. Stanley Williams  
Department of Chemistry  
University of California  
Los Angeles, California 90024

Dr. R. P. Messmer  
Materials Characterization Lab.  
General Electric Company  
Schenectady, New York 22217

Dr. Robert Gomer  
Department of Chemistry  
James Franck Institute  
5640 Ellis Avenue  
Chicago, Illinois 60637

Dr. Ronald Lee  
R301  
Naval Surface Weapons Center  
White Oak  
Silver Spring, Maryland 20910

Dr. Paul Schoen  
Code 6190  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. John T. Yates  
Department of Chemistry  
University of Pittsburgh  
Pittsburgh, Pennsylvania 15260

Dr. Richard Greene  
Code 5230  
Naval Research Laboratory  
Washington, D.C. 20375-5000

Dr. L. Kesmodel  
Department of Physics  
Indiana University  
Bloomington, Indiana 47403

Dr. K. C. Janda  
University of Pittsburgh  
Chemistry Building  
Pittsburg, PA 15260

Dr. E. A. Irene  
Department of Chemistry  
University of North Carolina  
Chapel Hill, North Carolina 27514

Dr. Adam Heller  
Bell Laboratories  
Murray Hill, New Jersey 07974

Dr. Martin Fleischmann  
Department of Chemistry  
University of Southampton  
Southampton SO9 5NH  
UNITED KINGDOM

Dr. H. Tachikawa  
Chemistry Department  
Jackson State University  
Jackson, Mississippi 39217

Dr. John W. Wilkins  
Cornell University  
Laboratory of Atomic and  
Solid State Physics  
Ithaca, New York 14853

ABSTRACTS DISTRIBUTION LIST, 056/625/629

Dr. R. G. Wallis  
Department of Physics  
University of California  
Irvine, California 92664

Dr. D. Ramaker  
Chemistry Department  
George Washington University  
Washington, D.C. 20052

Dr. J. C. Hemminger  
Chemistry Department  
University of California  
Irvine, California 92717

Dr. T. F. George  
Chemistry Department  
University of Rochester  
Rochester, New York 14627

Dr. G. Rubloff  
IBM  
Thomas J. Watson Research Center  
P.O. Box 218  
Yorktown Heights, New York 10598

Dr. Horia Metiu  
Chemistry Department  
University of California  
Santa Barbara, California 93106

Dr. W. Goddard  
Department of Chemistry and Chemical  
Engineering  
California Institute of Technology  
Pasadena, California 91125

Dr. P. Hansma  
Department of Physics  
University of California  
Santa Barbara, California 93106

Dr. J. Baldeschwieler  
Department of Chemistry and  
Chemical Engineering  
California Institute of Technology  
Pasadena, California 91125

Dr. J. T. Keiser  
Department of Chemistry  
University of Richmond  
Richmond, Virginia 23173

Dr. R. W. Plummer  
Department of Physics  
University of Pennsylvania  
Philadelphia, Pennsylvania 19104

Dr. E. Yeager  
Department of Chemistry  
Case Western Reserve University  
Cleveland, Ohio 44106

Dr. N. Winograd  
Department of Chemistry  
Pennsylvania State University  
University Park, Pennsylvania 16802

Dr. Roald Hoffmann  
Department of Chemistry  
Cornell University  
Ithaca, New York 14853

Dr. A. Steckl  
Department of Electrical and  
Systems Engineering  
Rensselaer Polytechnic Institute  
Troy, New York 12181

Dr. G.H. Morrison  
Department of Chemistry  
Cornell University  
Ithaca, New York 14853